

Quick Review of High School Mathematics

Ray Chien

July 27, 2026

Acknowledgements

I would like to gratefully acknowledge [Zhenverse](#)'s generous assistance. Zhenverse introduced LaTeX, and some valuable theorems to me, and taught me about appropriate mathematical statements. Of course, if there is any mistake in this work, it's all on me.

Contents

1	Book 1	2
1.1	Distance of a Point From a Line	2
1.2	Distance Between Two Parallel Lines	2
1.3	Equation of a Circle Given The Endpoints of a Diameter	3
1.4	Equations of a Tangent Line or a Chord of Contact on A Circle	3
1.5	Lagrange Interpolation	3
2	Book 2	3
2.1	Series	3
2.2	Percentiles	3
2.3	Variance and Standard Deviation	4
2.4	Correlation Coefficient	4
2.5	Line of Best Fit	4
2.6	Pascal's Principle	4
2.7	Combination With Repetition	4
2.8	Derangement	4
2.9	Sine Rule	4
2.10	Cosine Rule	4
3	Book 3	5
3.1	Projection Vector And Its Length	5
3.2	Triangle Inequality	5
3.3	Cauchy-Schwarz Inequality	5
3.4	Two-Dimensional Determinant	5
3.5	Cramer's Rule (2×2)	5

4	Book 4	6
4.1	Theorem of Three Perpendiculars	6
4.2	Cross Product	6
4.3	Volume of Parallelepipeds	6
4.4	Laplace Expansion	6
4.5	Point-Normal Form	7
4.6	Distance of a Point From a Plane	7
4.7	Distance Between Two Parallel Planes	7
4.8	Lines in Space	7
4.9	Conditional Probability	7
4.10	Independent Events	7
4.11	Bayes' Theorem	7
4.12	Operational Properties of Matrices	8
4.13	Inverse Matrix	8
4.14	Scaling Matrix	8
4.15	Shear Matrix	8
4.16	Rotation Matrix	8
4.17	Reflection Matrix	8
4.18	Transition Matrix	8
5	Additional Tricks	8
5.1	Centroids of Triangles	8
5.2	Circumcenters of Triangles	9
5.3	Incenters of Triangles	9
5.4	Orthocenters of Triangles	9
5.5	The Pairing Method	9
5.6	Stewart's Theorem	9
5.7	Vectors and Areas about a Point inside a Triangle	9
5.8	Areas of Triangles in Terms of Circumradius	10
5.9	Shoelace Formula	10
5.10	Brahmagupta–Fibonacci Identity	10

This work is licensed under a [Creative Commons](#)
[“Attribution-NonCommercial-ShareAlike 4.0](#)
[International”](#) license.



1 Book 1

1.1 Distance of a Point From a Line

$$\frac{|ax_0 + by_0 + c|}{\sqrt{a^2 + b^2}}$$

1.2 Distance Between Two Parallel Lines

$$\frac{|c_1 - c_2|}{\sqrt{a^2 + b^2}}$$

1.3 Equation of a Circle Given The Endpoints of a Diameter

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$$

1.4 Equations of a Tangent Line or a Chord of Contact on A Circle

$$(x - h)(x_0 - h) + (y - k)(y_0 - k) = r^2$$

$$x_0x + y_0y + d\frac{x + x_0}{2} + e\frac{y + y_0}{2} + f = 0$$

1.5 Lagrange Interpolation

$$\sum_{j=0}^n f(x_j) \left(\prod_{\substack{i=0 \\ i \neq j}}^n \frac{x - x_i}{x_j - x_i} \right)$$

2 Book 2

2.1 Series


$$\sum_{k=1}^n ar^{k-1} = \frac{a(1 - r^n)}{1 - r}$$

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{k=1}^n k^3 = \left[\frac{n(n+1)}{2} \right]^2$$

2.2 Percentiles



2.3 Variance and Standard Deviation

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2 = \frac{1}{n} \sum_{i=1}^n x_i^2 - \mu^2$$
$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2} = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - \mu^2}$$

2.4 Correlation Coefficient

$$r = \frac{1}{n} \sum_{i=1}^n x'_i y'_i = \frac{S_{xy}}{\sqrt{S_{xx}} \sqrt{S_{yy}}}$$

Note: The numerator is proportional to n , and the denominator is proportional to $\sqrt{n} \times \sqrt{n} = n$, so no extra multiplication or division by n is required.

2.5 Line of Best Fit

$$y - \mu_y = r \times \frac{\sigma_y}{\sigma_x} (x - \mu_x) = \frac{S_{xy}}{S_{xx}} (x - \mu_x)$$

2.6 Pascal's Principle

$$C_k^m = C_{k-1}^{m-1} + C_k^{m-1}$$

2.7 Combination With Repetition

$$H_k^n = C_k^{n+k-1}$$

2.8 Derangement

$$D_n = n! \sum_{k=0}^n \frac{(-1)^k}{k!}$$

2.9 Sine Rule

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$$

2.10 Cosine Rule

$$a^2 = b^2 + c^2 - 2bc \cos A$$

3 Book 3

3.1 Projection Vector And Its Length

Projection Vector:

$$\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \cdot \frac{\vec{b}}{|\vec{b}|}$$

Length of Projection Vector:

$$\frac{|\vec{a} \cdot \vec{b}|}{|\vec{b}|}$$

3.2 Triangle Inequality

$$||x| - |y|| \leq |x \pm y| \leq |x| + |y|$$

3.3 Cauchy-Schwarz Inequality

$$(a_1^2 + a_2^2)(b_1^2 + b_2^2) \geq (a_1b_1 + a_2b_2)^2$$

The equality holds if and only if $a_1b_2 = a_2b_1$

3.4 Two-Dimensional Determinant

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = - \begin{vmatrix} b & a \\ d & c \end{vmatrix}; \quad \begin{vmatrix} a & b \\ c & d \end{vmatrix} = - \begin{vmatrix} c & d \\ a & b \end{vmatrix}$$

$$\begin{vmatrix} ra & a \\ rc & c \end{vmatrix} = 0; \quad \begin{vmatrix} ra & rb \\ a & b \end{vmatrix} = 0$$

$$\begin{vmatrix} ra & b \\ rc & d \end{vmatrix} = r \begin{vmatrix} a & b \\ c & d \end{vmatrix}; \quad \begin{vmatrix} ra & rb \\ c & d \end{vmatrix} = r \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

$$\begin{vmatrix} a & ka+b \\ c & kc+d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}; \quad \begin{vmatrix} a & b \\ ka+c & kb+d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

$$\begin{vmatrix} a+x & b \\ c+y & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} x & b \\ y & d \end{vmatrix}; \quad \begin{vmatrix} a+x & b+y \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} x & y \\ c & d \end{vmatrix}$$

3.5 Cramer's Rule (2×2)

$$\begin{cases} a_1x + b_1y = c_1 \\ a_2x + b_2y = c_2 \end{cases}$$


$$\Delta = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}, \quad \Delta_x = \begin{vmatrix} c_1 & b_1 \\ c_2 & b_2 \end{vmatrix}, \quad \Delta_y = \begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix}$$

Condition	Solution	Geometric Meaning
$\Delta \neq 0$	Unique $(x = \frac{\Delta_x}{\Delta}, y = \frac{\Delta_y}{\Delta})$	Intersecting lines
$\Delta = 0, (\Delta_x \neq 0 \vee \Delta_y \neq 0)$	None	Parallel distinct lines
$\Delta = \Delta_x = \Delta_y = 0$	Infinite	Coincident lines

4 Book 4

4.1 Theorem of Three Perpendiculars

Let line AB be perpendicular to plane E at point B . If on plane E the line BC is perpendicular to line L at C , then line AC is perpendicular to line L at point C .



4.2 Cross Product

$$(a_1, a_2, a_3) \times (b_1, b_2, b_3) = \left(\begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix}, \begin{vmatrix} a_3 & a_1 \\ b_3 & b_1 \end{vmatrix}, \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \right)$$

4.3 Volume of Parallelepipeds

$$|\vec{a} \cdot \vec{b} \times \vec{c}| = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

4.4 Laplace Expansion

$$\begin{vmatrix} + & - & + \\ - & + & - \\ + & - & + \end{vmatrix}$$

4.5 Point-Normal Form

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

4.6 Distance of a Point From a Plane

$$\frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

4.7 Distance Between Two Parallel Planes

$$\frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}$$

4.8 Lines in Space

$$\begin{cases} x = x_0 + at \\ y = y_0 + bt \\ z = z_0 + ct \end{cases}, t \in \mathbb{R}$$
$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

4.9 Conditional Probability

$$P(B|A) = \frac{n(A \cap B)}{n(A)} = \frac{P(A \cap B)}{P(A)}$$

$$P(A \cap B) = P(B)P(A|B) = P(A)P(B|A)$$

$$P(B) = P(A' \cap B) + P(A \cap B)$$

4.10 Independent Events

$$P(A \cap B) = P(A)P(B)$$

If the events A, B are independent, then A', B are independent; A, B' are independent; and A', B' are independent.

4.11 Bayes' Theorem

$$P(A|B) = \frac{P(A)P(B|A)}{P(B)} = \frac{P(A)P(B|A)}{P(A)P(B|A) + P(A')P(B|A')}$$

4.12 Operational Properties of Matrices

$$(AB)C = A(BC)$$

$$A(B + C) = AB + AC$$

$$(A + B)C = AC + BC$$

4.13 Inverse Matrix

$$A^{-1} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

4.14 Scaling Matrix

$$\begin{bmatrix} h & 0 \\ 0 & k \end{bmatrix}$$

4.15 Shear Matrix

$$\begin{bmatrix} 1 & h \\ 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix}$$

4.16 Rotation Matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

4.17 Reflection Matrix

$$\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$$

4.18 Transition Matrix

$$\begin{bmatrix} a & b \\ 1 - a & 1 - b \end{bmatrix}$$

where $0 \leq a, b \leq 1$

5 Additional Tricks

5.1 Centroids of Triangles

$$\overrightarrow{DG} = \frac{1}{3}\overrightarrow{DA} + \frac{1}{3}\overrightarrow{DB} + \frac{1}{3}\overrightarrow{DC}$$

5.2 Circumcenters of Triangles

$$\overrightarrow{AB} \cdot \overrightarrow{AO} = \frac{1}{2} \overline{AB}^2$$

5.3 Incenters of Triangles

$$\overrightarrow{DI} = \frac{\overline{BC}}{\overline{AB} + \overline{BC} + \overline{CA}} \overrightarrow{DA} + \frac{\overline{CA}}{\overline{AB} + \overline{BC} + \overline{CA}} \overrightarrow{DB} + \frac{\overline{AB}}{\overline{AB} + \overline{BC} + \overline{CA}} \overrightarrow{DC}$$

5.4 Orthocenters of Triangles


$$\overrightarrow{AB} \cdot \overrightarrow{AC} = \overrightarrow{AB} \cdot \overrightarrow{AH} = \overrightarrow{AC} \cdot \overrightarrow{AH}$$

5.5 The Pairing Method

While dividing some labeled items into groups of two, pick the partners for the unpaired items. For example, the numbers of ways to pair 5 people and one person left is $C_1^5 \times 3 \times 1 = 15$.

5.6 Stewart's Theorem

$$z^2 = \frac{n}{m+n} x^2 + \frac{m}{m+n} y^2 - mn$$




When the dividing line is the bisector, $z^2 = xy - mn$.

When it is the median, $2(z^2 + n^2) = x^2 + y^2$.

5.7 Vectors and Areas about a Point inside a Triangle

$$S_A \vec{PA} + S_B \vec{PB} + S_C \vec{PC} = \vec{0}$$



5.8 Areas of Triangles in Terms of Circumradius

$$Area = \frac{abc}{4R}$$

5.9 Shoelace Formula

$$Area = \frac{1}{2} \left| \begin{vmatrix} x_1 & x_2 & \dots & x_k & x_1 \\ y_1 & y_2 & \dots & y_k & y_1 \end{vmatrix} \right|$$

5.10 Brahmagupta–Fibonacci Identity

$$(a^2 + b^2)(c^2 + d^2) = (ac + bd)^2 + (ad - bc)^2$$

The identity can be a trivial proof of the two-dimensional Cauchy–Schwarz inequality.